

# **IPRAL 250**

## **LIQUID FUNGICIDE**

**GROUP 2 FUNGICIDE**

**Active Constituent:** 250g/L IPRADIONE  
332g/L LIQUID HYDROCARBONS

**Poison Schedule:** S5 (CAUTION)

**APVMA Approval No:** 62283



Eligible containers  
10L  
5 L

For control of fungal diseases on various crops and situations as specified in the Directions for Use Table

**Pack Size:**

5L ( 4 x 5L per carton)  
10L

**Dangerous Goods Class:**

Not classified as a dangerous good under the Australian Code for Transport and Storage of Dangerous Goods in Australia.

---

**- DIRECTIONS FOR USE TABLE STARTS ON NEXT PAGE -**

# IPRAL 250

TREE CROPS/ VINES					
Rate				Critical Comments	
<b>RATE</b> In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Special Instructions for Tree Crops/Vines section.				<b>CRITICAL COMMENTS</b> For all uses in this table: Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Refer to the Special Instructions for Tree Crops/Vines section.	
Crop	Disease	State	Rate	WHP (days)	Critical Comments
Almonds	Blossom Blight, Bown Rot ( <i>Monilinia</i> spp., <i>Sclerotinia</i> spp.)	All States	100mL/100L water	Nil	Apply first at full bloom and, if conditions are favourable for disease development, up to two subsequent applications can be made: at petal fall and up to four weeks after petal fall.
Boysenberries	Grey Mould ( <i>Botrytis cinerea</i> )		200mL/100L water	1 day (H)	Spray at 10% blossom and full bloom. For fruit protection apply at 2 to 3 weeks pre-harvest.
Grapes	Grey Mould ( <i>Botrytis cinerea</i> )			7 days (H)	Good crop hygiene will aid in the control of disease. This use is subject to an Avcare fungicide resistance management strategy: 1. If three or fewer bunch rot sprays are applied in a season use only one spray per season containing Ipral (or other Group 2 Fungicide). If four or more bunch rot sprays are applied in a season use no more than two sprays containing Group 2 fungicides, unless tank mixed with a registered multi-site (Group Y) fungicide. 2. Late season fungicide treatments should be applied before botrytis infection reaches unacceptably high levels in the vineyard. 3. DO NOT apply more than two consecutive sprays from the same fungicide group, including from the end of one season to the next.
Kiwifruit	Botrytis Blight ( <i>Botrytis</i> spp.)	NSW, Vic, WA only			Apply the spray to vines every 10 to 14 days ensuring that all fruit is thoroughly wet. Apply 3 applications at 10 to 14 day intervals from 10% bloom to petal fall for protection of flowers and young fruit. Apply a further 2 applications of Ipral to control late season Botrytis.

# IPRAL 250

Crop	Disease	State	Rate	WHP (days)	Critical Comments
Macadamias	Botrytis Blight ( <i>Botrytis</i> spp.)	All States	100mL/ 100L water	Nil	Apply as a thorough cover spray to flower racemes when they open. A follow up spray may be needed one week later if wet conditions persist during flowering. Remove nuts under trees prior to spraying.
Mandarins (non-bearing)	Alternaria Leaf Spot (Brown Spot) ( <i>Alternaria alternata</i> )	Qld, WA, NT only	200mL/ 100L water		Apply to non-bearing trees of Murcott variety monthly from first flush in Spring until flushing ceases in the Autumn. Reduce intervals to fortnightly during periods of wet weather.
Passion fruit	Alternata Spot (Brown Spot) ( <i>Alternaria</i> spp., <i>Alternaria passiflorae</i> )	Qld, NSW, WA, NT only		7 days (H)	This use is subject to an Avcare fungicide resistance management strategy: 1. Maintain a protective cover with protectant fungicide such as Mancozeb. 2. Limit the use of Ipral to strategic periods. i.e. before, during and after extended wet periods. 3. Always tank mix Ipral with a protectant such as Mancozeb. 4. DO NOT apply more than four Ipral (or other Group 2 fungicide) sprays in a season.
Raspberries	Grey Mould ( <i>Botrytis cinerea</i> )	All States	200mL/ 100L water	1 day (H)	Spray at 10% blossom and full bloom. For fruit protection, apply at 2 to 3 weeks pre-harvest.

# IPRAL 250

Crop	Disease	State	Rate	WHP (days)	Critical Comments
Stone fruit: Apricots, Cherries, Nectarines, Peaches, Plums	Orchard spraying Blossom Blight ( <i>Monilinia fructicola</i> , <i>Monilinia laxa</i> )  Brown Rot ( <i>Monilinia fructicola</i> , <i>Monilinia laxa</i> )	Qld, NSW, Vic, Tas, SA, WA only	100 to 150mL/ 100L water	Nil	For control of Blossom Blight, spray at 10% blossom, full bloom and petal/shuck fall. For control of subsequent Brown Rot in fruit, spray at 3 weeks and 1 week preharvest. Use the higher rate under severe conditions of challenge, or for single applications of Ipral in the spray program. This use is subject to an Avcare fungicide resistance management strategy: 1. DO NOT apply more than 2 consecutive sprays of Ipral (or other Group 2 fungicides). 2. A post-harvest treatment should also be counted as an application. 3. The last blossom blight spray and the first pre-harvest brown rot spray should be regarded as consecutive applications. 4. The spray program should be considered and the strategy applied on a whole-orchard basis.
Youngberries	Grey Mould ( <i>Botrytis cinerea</i> )	All States	200mL/ 100L water	1 day (H)	Spray at 10% and full bloom. For fruit protection, apply at 2 to 3 weeks pre-harvest.
Strawberries	Grey Mould ( <i>Botrytis cinerea</i> )	All States	2.0L/ha where spray volume is less than 1000L/ha OR 200mL/100L water where spray volume equals or exceeds 1000L/ha	1 day (H)	This use is subject to an Avcare fungicide resistance management strategy: 1. Apply a program of protectant fungicides during flowering. If conditions favour disease development during this period use Ipral. 2. DO NOT apply more than two successive sprays of Ipral (or other Group 2 Fungicide).
Berries (See Tree Crops/Vines for boysenberries, raspberries and youngberries)					

# IPRAL 250

Vegetables					
Crop	Disease	State	Rate	WHP (days)	Critical Comments
Celery	Sclerotinia Rot (Pink Rot) ( <i>Sclerotinia sclerotiorum</i> )	All States	2.0L/ha where spray volume is less than 1000L/ha OR 200mL/100L water where spray volume equals or exceeds 1000L/ha	1 day (H)	Commence spraying 1 to 2 weeks post-transplanting and then every 2 to 3 weeks. Use only five sprays.
Lettuces	Sclerotinia Rot ( <i>drop</i> ) ( <i>Sclerotinia sclerotiorum</i> , <i>Sclerotinia minor</i> )			7 days (H)	
	Grey Mould ( <i>Botrytis</i> spp.)	Tas, WA only			Spray should be directed to the stems at ground level and to the underside of lower leaves. This use is subject to an Avcare fungicide resistance management strategy: <ol style="list-style-type: none"> <li>1. Apply Ipral as a seedling drench soon after emergence.</li> <li>2. Apply a protectant fungicide as a high volume foliar spray before planting out, then Ipral immediately after planting.</li> <li>3. Maintain cover with protectant fungicide sprays at 7-10 day intervals</li> <li>4. If weather conditions favour <i>Botrytis</i> infection, tank mix the protectant with Ipral.</li> <li>5. Do not apply Ipral (or other Group 2 fungicides) more than four times per season, irrespective of the target disease.</li> </ol>

# IPRAL 250

Crop	Disease	State	Rate	WHP (days)	Critical Comments
Potatoes	Sclerotinia Rot ( <i>Sclerotinia sclerotiorum</i> )	All States	1.0 to 2.0L/ha where spray volume is less than 1000L/ha OR 100 to 200mL/100L water where spray volume equals or exceeds 1000L/ha	Nil	Apply 2 sprays, once immediately before and once immediately after hilling-up. For most effective treatment, concentrate the spray at the base of the stems and surrounding soil surface, where the fungus is active. Use higher rate where disease is severe.
	Target Spot (Early Blight) ( <i>Alternaria solani</i> )				Ensure thorough coverage to the whole plant. Treatment is generally not required until after flowering. Use the higher rate where disease is severe. This use is subject to an Avcare fungicide resistance management strategy: <ol style="list-style-type: none"> <li>1. Limit the use of Ipral to periods when conditions favour disease development.</li> <li>2. DO NOT apply more than four Ipral (or other Group 2 fungicide) sprays in one season.</li> <li>3. Apply no more than two consecutive sprays of a Group 2 fungicide.</li> </ol>
	Hypocotyl Rot (Black Scurf) ( <i>Rhizoctonia solani</i> )		800mL/tonne seed material		Ipral will protect emerging shoots from hypocotyl rot, improving overall germination. Ipral may also reduce occurrence of black scurf on the harvested potatoes. Ensure good coverage of seed material and planting furrow. This can be achieved by applying Ipral as a fine spray to the seed at the time of planting using spray equipment mounted on the planter, and nozzles located at three points on each planter row to ensure uniform coating of the seed. DO NOT plant into waterlogged soil. A minimum water volume of 80L/tonne seed should be used.

# IPRAL 250

Crop	Disease	State	Rate	WHP (days)	Critical Comments
Tomatoes	Sclerotinia Rot ( <i>Sclerotinia sclerotiorum</i> )	Qld, NSW, Tas, SA, WA only	2.0L/ha where spray volume is less than 1000L/ha OR 200mL/100L water where spray volume equals or exceeds 1000L/ha	7 days (H)	Spray at 14-day intervals from transplanting and throughout the period of disease pressure.
	Grey Mould ( <i>Botrytis cinerea</i> )	All States			Commence spraying 3 to 4 weeks after transplanting or at the onset of disease. Repeat treatment at 14-day intervals or when conditions favour spread of the disease, i.e. at trimming or deleafing. This use is subject to an Avcare fungicide resistance management strategy: 1. Alternate or tank mix Ipral with a protectant such as chlorothalonil. Avoid applying two Ipral (or other Group 2 fungicide) sprays in succession, unless tank mixed with a protectant. 2. Do Not apply more than four Ipral (or other Group 2 fungicide) sprays in a season.
	Target Spot ( <i>Early Blight</i> ) ( <i>Alternaria solani</i> )	Qld, Tas, WA, NT only			Commence spraying 1 week posttransplanting. Use adequate water to give thorough coverage of the plants. Use high volume spray equipment. This use is subject to an Avcare fungicide resistance management strategy: 1. Limit the use of Ipral to period when conditions favour disease development. 2. Do Not apply more than four Ipral (or other Group 2 fungicide) sprays in one season. Apply no more than two consecutive sprays of a Group 2 fungicide.

# IPRAL 250

FIELD CROPS					
Crop	Disease	State	Rate	WHP (days)	Critical Comments
Canola	Sclerotinia Stem Rot (Sclerotinia sclerotiorum)	All States	2.0L/ha	6 weeks (H,G)	Apply at 20 to 50% flowering. Apply as a preventative spray before disease infection is anticipated. Good coverage is essential. Aerial application: Apply using a minimum water volume of 45L/ha. Ground application: Apply using a minimum water volume of 100L/ha.
Lucerne	Lucerne Leaf Spot ( <i>Stemphylium botryosum</i> )	Qld, WA only	500mL to 1.0L/ha where spray volume is less than 1000L/ha OR 50-100mL/100L water where spray volume equals or exceeds 1000L/ha	7 days (G)	Spray every 10 to 14 days when cool, damp weather favours the disease. Use the higher rate under conditions of high disease pressure.
	Leptosphaerulina Leaf Spot ( <i>Leptosphaerulina trifolii</i> )				Apply in at least 300L water/ha every 10 to 14 days when cool, damp weather favours the disease. Use the higher rate under conditions of high disease pressure.
Peanuts	Sclerotinia Rot (Sclerotinia sclerotiorum, Sclerotinia minor)		2.0L/ha OR 440mL/100L water (spot application)	12 days (H)	Apply when disease first appears. Repeat if necessary. Use a high water volume to ensure good coverage of foliage and stem at ground. Do not mix Ipral with a foliar fungicide due to the different target positions on the plant.
Soybeans	Black Leaf Blight ( <i>Arkoala nigra</i> )	NSW, WA only	2.0L/200 to 400L water/ha	7 weeks (H)	If disease is present on leaves apply an initial spray at early pod set (pods approximately 5mm long). An additional spray 14 days later may be required if wet seasonal conditions prevail.
<b>NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.</b>					

# IPRAL 250

WITHHOLDING PERIODS (WHP): (H = HARVEST, G = GRAZING)

Almonds, macadamias, mandarins, potatoes, stone fruit: NOT REQUIRED WHEN USED AS DIRECTED.

Boysenberries, celery, raspberries, strawberries, youngberries: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

Grapes, kiwifruit, lettuce, tomatoes and passion fruit: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

Peanuts: DO NOT HARVEST FOR 12 DAYS AFTER APPLICATION.

Canola: DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION. DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 6 WEEKS AFTER APPLICATION.

Soybeans: DO NOT HARVEST FOR 7 WEEKS AFTER APPLICATION.

Lucerne: DO NOT GRAZE OR CUT FOR STOCK FOOD WITHIN 7 DAYS OF TREATMENT.

## GENERAL INSTRUCTIONS

GROUP	2	FUNGICIDE
-------	---	-----------

### Fungicide Resistance Warning

Ipral Fungicide is a member of the dicarboximide group of fungicides. For fungicide resistance management the product is a Group 2 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 2 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 2 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Colin Campbell (Chemicals) Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

### Resistance Management

Resistant strains of fungi can develop to this and other fungicides. To reduce the possibility of this occurrence, and where alternatives are available, rotate to use products with as many different modes of action as possible. Where specific resistance management strategies are established these are detailed in the Critical Comments.

### Export of Treated Produce

Growers should note that MRLs or import tolerances may not exist in all markets for produce treated with Ipral. If you are growing produce for export, please check with Colin Campbell (Chemicals) Pty Ltd, for the latest information on MRLs and import tolerances BEFORE using Ipral.

### Mixing

Note: Ipral may be unstable in conditions where the pH is 7 or higher. It is therefore essential to check the pH of the spray solution before adding Ipral. A suitable registered buffering agent may have to be added to bring the pH down below 7. Shake well before use. Add half the required water volume to the spray tank or vat with the agitation mechanism operating. Add the required volume of this product and then add additional water to the volume required.

### Application

Good disease control requires even thorough coverage of the target area. Application should be made using appropriate spray equipment and sufficient water to provide adequate penetration and coverage. Equipment settings and water volume may need to vary, depending on the growth stage of the crop. High pressure, prolonged and vigorous agitation particularly in conjunction with a high concentration of Ipral in the spray tank may reduce the suspension properties of Ipral, resulting in a scum forming on the surface or sediment forming on the filters.

If the agitation system cannot be adjusted, or concentration reduced to overcome this problem it is recommended that Ippon Aquaflo be used, where registered.

# IPRAL 250

## Special Instructions for Tree Crops/Vines

### Dilute Spraying

- Use a sprayer designed to apply high spray volumes, up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient spray solution to cover the crop to the point of run-off. Avoid excessive run-off.
- The required spray volume to achieve point of run off may be determined by applying different test volumes, using different settings on the sprayer, or from industry guidelines or other expert advice.
- Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run-off.
- The required dilute spray volume to achieve point of run off will change and the sprayer set up and operation may also need to be changed, as the crop grows.

### Concentrate Spraying

- Use a sprayer designed and set up to concentrate spraying (that is a sprayer which applies spray volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen spray volume.
- Determine an appropriate dilute spray volume (See Dilute spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:

#### EXAMPLE ONLY

1. Dilute spray volume as determined above: For example 1500L/ha
  2. Your chosen concentrate spray volume: For example 500L/ha
  3. The concentration factor in this example is:  $3 \times$  (i.e.  $1500L \div 500L=3$ )
  4. If the dilute label rate is 10mL/100L, then the concentrate rate becomes  $3 \times 10$ , that is 30mL of product per 100L water for concentrate spraying.
- The chosen spray volume, amount of product per 100L of water, and the sprayer set up and operation may need to be changed as the crop grows.
  - For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

### Compatibility

Ipral is compatible with the following products:

Aliette WG (see NOTE below), azinphos-methyl, benomyl, Bugmaster Flowable, chlorfenvinphos, chlorpyrifos (500g/L EC), demeton-S-methyl, Dithane M45, Thiodan EC (endosulfan), fenarimol, Kelthane, Kocide (Warning: Do not mix Ipral with Kocide for use on potatoes), Larvin 375, Maldison 500, Marlin, metalaxyl, methamidophos, methyl parathion, pirimicarb, propargite, triadimenol. When tank mixing products the order of mixing is determined by formulation type. As a guide the following mixing sequence is recommended:

1. Wettable powders
2. Suspension concentrates
3. Water Dispersible Granules
4. Suspo-emulsions (e.g Ipral)
5. Soluble powders
6. Solutions
7. Emulsifiable concentrates
8. Soluble concentrates
9. Wetting agents and oils

# **IPRAL 250**

With any mixture, thoroughly agitate immediately before applying. It is not recommended to mix this product with more than one of the above chemicals in the tank. The use of a surfactant or spray oil is not recommended with Ipral as it may result in crop damage to sensitive plants. DO NOT mix with fertilisers. Mixtures with some fertilisers, e.g urea, may cause foliar damage. Note: Mixing Ipral with Aliette WG may result in some settling out. As formulation of other manufacturers products are beyond the control of Colin Campbell (Chemicals) Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS AND ORGANISMS:** DO NOT apply the product under weather conditions, or from spraying equipment, which could be expected to cause spray drift onto adjacent crops, croplands, pastures, livestock, natural or impounded lakes, dams or other waterways.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:** Toxic to aquatic organisms. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

**STORAGE AND DISPOSAL:** Store in a closed, original container in a cool, secure, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Protect from frost. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. 110 L containers only: Empty container by pumping through the dry-break connection system. Do not attempt to unscrew the valve or breach the locked filling point. Do not contaminate the container with water or other foreign material. Ensure that the coupler, pump, meter and hoses are disconnected, triple rinsed with clean water and drained after each use. Contact point of purchase to arrange return or collection of empty containers.

**SAFETY DIRECTIONS:** Avoid contact with eyes and skin, and avoid inhalation of vapour. Wear suitable protective clothing, gloves and goggles. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre (Phone 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.

## **MATERIAL SAFETY DATA SHEET**

Additional information is contained in the Material Safety Data Sheet. A copy can be obtained from the website:

### **CONDITIONS OF SALE**

Agricultural, horticultural and pastoral preparations in their application involve varying factors such as differing conditions, soil, climate and methods of application over which the vendor does not have control, Whilst Colin Campbell (Chemicals) Pty. Ltd believes that all goods sold by it are true to label and are effective and safe for purpose indicated the company and the seller hereby expressly negate and exclude any express or implied condition, statement or warranty, statutory or otherwise, as to quality or fitness of any goods sold for any purpose or purposes whatsoever except such warranties and conditions, if any, as are implied by the Trade Practices Act 1974 (Commonwealth). The company and the seller accept no responsibility for any loss, harm or damage whatsoever suffered from the use of such goods for any purpose or purposes irrespective of whether or not the buyer was acting in reliance upon the advice recommendation or representation of the seller or any representative agent of employee of the company as to such use except in respect of breaches of conditions and warranties, if any, implied by the Trade practices Act and in respect of such breaches the liability of the company and the seller shall be limited to the replacement of the goods or the supply of equivalent goods, or the payment of the cost of replacing the goods.

APVMA Approval No: 62283

®Ipral is a registered trademark of Colin Campbell (Chemicals) Pty Ltd