



# PROTAK

GROUP **C** FUNGICIDE

**Active Ingredient:** 450g/L Prochloraz  
Solvent: 390g/L Xylene

**Poison Schedule:** S6

**Resistant Grouping:** Group C

**Dangerous Goods Class:** 3

**Registered for use In:** N.S.W., VIC., S.A., QLD., N.T., W.A.

**Available In:** 1L (12 x 1L per carton)

## DIRECTIONS FOR USE

CROP	DISEASE	STATES	RATE	CRITICAL COMMENTS
AVOCADO	Anthrachnose Stem End Rot	Qld, NSW, NT, WA only	55mL/100L water	Spray fruit for 30 seconds. DO NOT use on Avocado cultivar Rincon.
BANANA	Anthrachnose (Black-End)			
MANGO	Anthrachnose ( <i>Colletotrichum gloeosporiodes</i> )			
	Alternaria Rot ( <i>Alternaria alternata</i> )	Qld only		
PAWPAW	Anthrachnose, Stem-end	Qld, NT, NSW, WA, only		Spray fruit for 1 minute.
PINEAPPLES	Water Blister ( <i>Ceratocystis paradoxa</i> )	Qld only		Dip fruit for 1 minute. As the dip volume declines top-up using a concentration of 85ml per 100L water.
SUGAR CANE	Pineapple disease	Qld, NSW, NT, only	40mL/200L water	Apply as a dip or spray to sets at planting. Ensure thorough coverage at all set ends.
TURF	Dollar Spot ( <i>Sclerotinia homeocarpa</i> )	Qld, NSW, Vic, SA, WA, NT only	65mL/50 to 100L water/100m <sup>2</sup>	Apply in a monthly spray programme when disease first appears. Spray evenly to ensure thorough coverage.



### **General Instructions**

**Mixing:** Add the required amount of product to the water in the spray tank/reservoir and stir thoroughly to ensure complete mixing. Add anti-foaming agent in situations where foaming of the dip solution could be a problem.

**Application - Post-Harvest:** Dip Pineapples as directed.

For other crops, until a topping-up recommendation is available, this product should be used in non-recirculating spray systems only. Treat fruit as soon as possible after harvest at shed temperature. Thorough coverage is essential. Adjust spraying equipment to ensure exposure of fruit for the periods recommended in the Critical Comments above. DO NOT use this product as a foliar spray on fruit as it may cause burning of foliage.

### **Environmental Information**

DO NOT graze stock on treated turf or feed grass clippings from treated turf to livestock.

Harmful to Fish. Do not contaminate ponds, streams, rivers, dams, drains or waterways with this product or used containers.

### **Storage and Disposal**

Store in the closed original container in a well ventilated area, as cool as possible. Do not store with fertilizers, pesticides and seeds. Do not store for prolonged periods in direct sunlight. Triple rinse containers adding rinsings to the spray tank, or dispose of with empty container and any excess spray. Puncture top, bottom and sides of empty container then crush and bury in an approved landfill or bury under at least 500mm of soil in a non-crop, non-pasture area away from water sources or homes.

### **Safety Directions**

Product is poisonous if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the product and preparing the spray wear elbow-length PVC gloves. If product on skin immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each days use wash gloves and contaminated clothing.

### **First Aid**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126. If swallowed, and if more than 15 minutes from a hospital induce vomiting, preferably using Ipecac Syrup APF. For further information refer to MSDS 9.

**APVMA Approval No: 45187**