



TRIDIM

GROUP **C** FUNGICIDE

Active Ingredient:	250g/L Triadimenol
Poison Schedule:	S6
Withholding period:	see below
Resistant Grouping:	Group C
Dangerous Goods Class:	Not classified as a dangerous good under the Australian Code For Transport and Storage of Dangerous Goods in Australia.
Registered for use In:	All States
Available In:	1L (10 x 1L per carton) 5L (4 x 5L per carton)



DIRECTIONS FOR USE

Restraint: Do not apply to turf earlier than 5 weeks after sowing.

CROP	DISEASE	STATE	RATE	CRITICAL COMMENTS
Cabbage Cauliflower Broccoli	Ring spot	All States	200 or 400 mL/ha plus a non-ionic wetter 200 mL/ha	Apply every 10 days. Start the program immediately following the first two consecutive days of rain after transplanting. Use higher rate under heavy disease pressure or when weather conditions favour development of the disease.
Cucurbits	Powdery mildew	All States	40 mL/100L or 400 mL/ha	Apply a non-DMI fungicide on a schedule commencing as soon as possible after emergence until fruit set. Then either tank mix or alternate Tridim with a non-DMI fungicide on a 5-10 day schedule. Use short spray intervals when disease pressure is higher plants are growing rapidly. Good coverage is essential for good results.
Grapevines	Powdery mildew	All States	10 mL/100L or 100 mL/ha	Apply thoroughly in the following 5 spray program: 1- when shoots are 10 cm long. 2- pre-flowering. 3- at flowering. 4- after fruit set. 5- before bunches close. Do not apply in volumes less than 250 L/ha. For spray volumes between 250 and 1000L/ha, use Tridim at 100 mL/ha. For volume greater than 1000L/ha, use the volume rate of 10 mL/100L.
Papaws	Powdery mildew	QLD, NSW, WA only	20 mL/100L or 200 mL/ha	Spray when powdery mildew infection begins or when conditions favour the disease. Repeat application every 14 days while conditons favour the disease. Ensure thorough coverage of foliage particularly the crown region.
Sugar cane	Pineapple disease	QLD, NSW only	20 mL/100L	Apply to sets by dipping or spraying. Ensure thorough wetting of cut ends.



CROP	DISEASE	STATE	RATE	CRITICAL COMMENTS
Turf	Dollar spot	QLD, NSW, VIC, SA, WA only	Preventive 30mL/100m ²	Apply thoroughly on a monthly programme starting immediately the disease appears.
			Curative 60mL/100m ²	Apply thoroughly under conditions of heavy disease pressure then treat as necessary.
	Black Helminthosporium	QLD, NSW, TAS only	60mL/100m ²	Use high volume equipment. Low volume may be used provided turf is irrigated thoroughly after spraying. Apply during period of turf growth on a monthly programme.
	Take – all patch	QLD, NSW, VIC, TAS, WA only		Preventive only after soil fumigation with methyl bromide. Apply thoroughly, using high volume equipment, 6-8 weeks after seeding. Repeat applications (every 8 weeks in the first two years) for continued protection.
Winter Fusarium	NSW, VIC, TAS only		Apply thoroughly on a monthly programme from end of March until danger of infection has passed.	

WITHHOLDING PERIODS FOR POST HARVEST USE:

CUCURBITS: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

CABBAGE, CAULIFLOWER, BROCCOLI, GRAPEVINES, PAPAWS: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

CABBAGE, CAULIFLOWER, BROCCOLI: DO NOT GRAZE ANY TREATED AREA, OR CUT FOR STOCK FOOD.

General Instructions

Tridim 250 EC is a broad spectrum, systemic, foliar fungicide that is quickly taken up by plants and translocated within the plant. Because of its systemic activity, Tridim is less affected by rainfall or irrigation than contact type fungicides.

Mixing

Half fill the spray tank or vat with clean water. Then add the required amount of Tridim with agitator going and top up the tank with water. Agitate thoroughly prior to and during spraying.

Protection of Wildlife, Fish, Crustacea and Environment

Harmful to fish. Do not contaminate streams, rivers or waterways with this product or used container.



Storage and Disposal

Store in the closed, original container in a well ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight. Triple or (preferably) pressure rinse container before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Safety Directions

Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length butyl rubber gloves and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

APVMA Approval No: 51580