

Syllit[®]

FUNGICIDE

GROUP M7 FUNGICIDE

Active Constituent: 400G/L DODINE
Poison Schedule: S6 (POISON)
APVMA Approval No: 54058



Eligible containers
10L

For the control of Black Spot in Apples and Pears, and Leaf Curl and Blossom Blight in Peaches and Nectarines.

Pack Size:
10L (single cubes)

Dangerous Goods Class:
9

DIRECTIONS FOR USE for all states				
Rate In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Mixing/Application section.				Critical Comments
Crop	Disease	Rate per 100L	WHP	
Apples	Black Spot (Venturia inaequalis)	80mL	5 days	PROTECTIVE PROGRAMME: Following a copper spray at green tip, apply at pink bud (apples) or white bud (pears) followed by 2 cover sprays 10 days apart, a further 2 cover sprays 10-14 days apart and a fifth cover spray after a further 21 days and then at 4-6 week intervals depending on weather and presence of infection CURATIVE PROGRAMME: Apply within 36 hours of the start of an infection period (e.g. after rain) which is favourable to development of black spot. *This fungicide is subject to resistance strategy. See General Instructions for details.
Pears	Black Spot (Venturia inaequalis)	120mL		
Peaches Nectarines	Peach leaf curl (Taphrina deformans)	100mL	-	Apply at bud swell, early bloom and petal fall. DO NOT apply after petal fall.
	Blossom Blight (Monilinia spp)	200mL		
NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORIZED UNDER APPROPRIATE LEGISLATION				

WITHHOLDING PERIOD:

APPLES AND PEARS: DO NOT HARVEST FOR 5 DAYS AFTER APPLICATION.

GRAZING: DO NOT CUT FODDER FOR STOCKFEED OR ALLOW LIVESTOCK TO GRAZE VEGETATION IN TREATED ORCHARDS.

GENERAL INSTRUCTIONS: This product possesses eradicant as well as protective properties. When applied within 36 hours after the start of an infection period, it penetrates the leaf surface to eradicate the black spot or scab infection.

DO NOT use on fruit at temperatures close to freezing point from pink stage on, as fruit russetting may occur. DO NOT use on apples or pears under conditions conducive to skin damage including russetting, lentical spotting and drip point burn. (Varieties such as Golden Delicious apples and Williams and Packham pears are particularly susceptible).

MIXING: Shake well before use. Add the necessary quantity to about half the required amount of water in the spray tank while the agitator is running. Then add the rest of the water and keep mixture agitated during use

APPLICATION

Dilute spraying

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate spraying

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (see Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way.

EXAMPLE ONLY

1. Dilute spray volume as determined above: for example 1500L/ha.
 2. Your chosen concentrate spray volume: for example 500L/ha.
 3. The concentration factor in this example is : 3 X (i.e. 1500L / 500L = 3)
 4. If the dilute label rate is 40mL/100L, then the concentrate rate becomes 3 x 40, that is 120mL/100L of concentrate spray.
- The chosen spray volume, amount of product per 100L of water, and the sprayer set up and operation may need to be changed as the crop grows.
 - For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practice.

FUNGICIDE RESISTANCE WARNING

GROUP	M7	FUNGICIDE
--------------	-----------	------------------

For fungicide resistance management Syllit is a Group M7 fungicide. Some naturally occurring individual fungi

resistant to Syllit and other Group M7 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Syllit and other Group M7 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use Colin Campbell (Chemicals) Pty. Ltd accepts no liability for any losses that may result from the failure of Syllit to control resistant fungi.

RESISTANCE STRATEGY:

Continual use of this product may result in the build up of strains of disease resistant to Syllit . To avoid resistance build up Colin Campbell (Chemicals) Pty Ltd recommend applying no more than 3 consecutive sprays of Syllit . If more than 3 sprays are required include a fungicide from DMI group before continuing with Syllit .

PRECAUTIONS:

Re-entry : Do not allow re-entry into treated areas for 2 days after spraying, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

COMPATIBILITY:

DO NOT mix with more than one product. DO NOT mix with lime sulphur, Bordeaux mixture, chlorbenzilate or alkaline materials. This product may be mixed with most commonly used insecticides, acaricides and fungicides. However, with some products, particularly emulsifiable concentrates and with some types of water, it may be desirable to add non-ionic wetting agent to the water before mixing Syllit or adding the other product. Particular combinations should be checked before preparing large amounts.

PROTECTION OF LIVESTOCK:

DO NOT graze cover crops in treated orchards or feed crop residues to livestock.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT:

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL:

Store in the closed, original container in a well-ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilizers. Triple rinse empty containers before disposal, and dispose of washings in a disposal pit away from plants and their roots. Wash sprayer thoroughly after use. Destroy empty container by breaking, crushing or puncturing. Bury the container at a depth of 500mm or more at a safe disposal site away from water sources or homes, or take to a dump that does not burn its refuse. DO NOT burn empty container or product.

SAFETY DIRECTIONS:

Harmful if inhaled or swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves, goggles and disposable mist mask. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing).

If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 03 4747000.

For further information, refer to the Material Safety Data Sheet (MSDS 74).

CONDITIONS OF SALE

Agricultural, horticultural and pastoral preparations in their application involve varying factors such as differing conditions, soil, climate and methods of application over which the vendor does not have control, Whilst Colin Campbell (Chemicals) Pty. Ltd believes that all goods sold by it are true to label and are effective and safe for purpose indicated the company and the seller hereby expressly negate and exclude any express or implied condition, statement or warranty, statutory or otherwise, as to quality or fitness of any goods sold for any purpose or purposes whatsoever except such warranties and conditions, if any, as are implied by the Trade Practices Act 1974 (Commonwealth). The company and the seller accept no responsibility for any loss, harm or damage whatsoever suffered from the use of such goods for any purpose or purposes irrespective of whether or not the buyer was acting in reliance upon the advice recommendation or representation of the seller or any representative agent of employee of the company as to such use except in respect of breaches of conditions and warranties, if any, implied by the Trade practices Act and in respect of such breaches the liability of the company and the seller shall be limited to the replacement of the goods or the supply of equivalent goods, or the payment of the cost of replacing the goods.

APVMA Approval No: 54058

®Syllit is a registered trademark of Agriphar SA